


CALIFORNIA FISH and GAME



"CONSERVATION THROUGH EDUCATION"

VOLUME 82

SUMMER 1996

NUMBER 3



California Fish and Game is published quarterly by the California Department of Fish and Game. It is a journal devoted to the conservation and understanding of California's flora and fauna. If its contents are reproduced elsewhere, the authors and the California Department of Fish and Game would appreciate being acknowledged.

Subscriptions may be obtained at the rate of \$15 (\$20 for foreign subscribers) per year by placing an order with the Editor, *California Fish and Game*, 4001 North Wilson Way, Stockton, California 95205-2486. Checks or money orders in U.S. dollars should be made out to *California Fish and Game*. Inquiries regarding subscriptions should be directed to the editors. Complimentary subscriptions are granted on an exchange basis.

California Department of Fish and Game employees may request complimentary subscriptions to the journal.

Please direct correspondence to:

David W. Kohlhorst or James J. Orsi
Co-Editors-in-Chief
California Fish and Game
4001 North Wilson Way
Stockton, California 95205-2486
e-mail: dkohlhor@delta.dfg.ca.gov
jorsi@delta.dfg.ca.gov



CALIFORNIA FISH AND GAME

VOLUME 82

SUMMER 1996

NUMBER 3



Published Quarterly by

STATE OF CALIFORNIA
THE RESOURCES AGENCY
DEPARTMENT OF FISH AND GAME
--LDA--

STATE OF CALIFORNIA
PETE WILSON, *Governor*

THE RESOURCES AGENCY
DOUGLAS P. WHEELER, *Secretary for Resources*

FISH AND GAME COMMISSION

Douglas B. McGeoghegan, *President*
Richard T. Thieriot, *Vice-president*
Frank D. Boren, *Member*
Robert R. Treanor, *Executive Director*

DEPARTMENT OF FISH AND GAME

JACQUELINE E. SCHAFER, *Director*
Vacant, *Chief Deputy Director*
R.A. Bernheimer, *Deputy Director*
Ryan Broddrick, *Deputy Director*
Kassandra Fletcher, *Deputy Director*
A. Petrovich Jr., *Deputy Director*

Perry L. Herrgesell, Ph.D., *Chief* Bay-Delta and Special Water Projects Division
John Turner, *Chief* Environmental Services Division
Timothy C. Farley, *Chief* Inland Fisheries Division
LB Boydstun, *Acting Chief* Marine Resources Division
Susan A. Cochrane, *Chief* Natural Heritage Division
Terry M. Mansfield, *Chief* Wildlife Management Division
DeWayne Johnston, *Chief* Wildlife Protection Division
Richard Elliott, *Regional Manager* Redding
Banky E. Curtis, *Regional Manager* Rancho Cordova
Brian Hunter, *Regional Manager* Yountville
George Nokes, *Regional Manager* Fresno
Patricia Wolf, *Acting Regional Manager* Long Beach
Pete Bontadelli, *Administrator* Oil Spill Prevention and Response

CALIFORNIA FISH AND GAME
EDITORIAL STAFF

James J. Orsi and David W. Kohlhorst, *Co-Editors-in-Chief* Bay-Delta
Walter Beer; John Brode; Ralph Carpenter; Arthur C. Knutson, Jr. ... Inland Fisheries
Vernon Bleich, Dan Yparraguirre Wildlife Management
Peter Kalvass, David Parker, Marija Vojkovich, Ronald Warner ... Marine Resources
Donald E. Stevens Bay-Delta
Liam Davis, Peter T. Phillips Environmental Services

CONTENTS

ARTICLES

- An Analysis of Commercial Passenger Fishing Vessel Fisheries
for Kelp Bass and Barred Sand Bass in the Southern
California Bight Milton S. Love, Andrew Brooks, and J. R. Raymond Ally 105
- Condition and Reproductive Performance of Female Mule Deer
in the Central Sierra Nevada T. J. Taylor 122

NOTES

- Southern Range Extension of the Harlequin Rockfish, *Sebastes
variegatus* (Scorpaenidae) James Wilder Orr and David C. Baker 133
- Distribution of Nesting Greater Sandhill Cranes in the South
Fork Pit River Valley, Modoc County, California Thomas A. Roberts, Rebecca Cull, and Anne Flannery 137
- A Red Abalone Tag Return after 16 Years at Liberty Ian K. Taniguchi and Peter L. Haaker 141
- First Freshwater Record of Pacific Lamprey, *Lamptera
tridentata*, from Baja California, Mexico Gorgonio Ruiz-Campos and Salvador Gonzalez-Guzman 144
- Past Occurrence of Eulachon, *Thaleichthys pacificus*, in
Streams Tributary to Humboldt Bay, California Mark R. Jennings 147

COVER

Male and Female Mule Deer

PUBLICATION NOTICE

A Bibliography of Feral, Stray, and Free-roaming Domestic Cats in Relation to Wildlife Conservation. 1994. Ronald M. Jurek. California Department of Fish and Game, Nongame Bird and Mammal Program Report No. 94-5. 24 pages. This is a comprehensive bibliography of 340 international references, mostly concerning effects of the domestic cat, *Felis catus*, on wildlife resource management and protection. The bibliography includes a subject index to aid the reader in identifying content of cited articles. The purpose of this bibliography is to provide an information source on the ecology and control of feral and free-roaming domestic cats and their damaging effects on wildlife. It is available free of charge from the author, c/o Bird and Mammal Conservation Program, California Department of Fish and Game, 1416 Ninth Street, Sacramento, California 95814-5509.

AN ANALYSIS OF COMMERCIAL PASSENGER FISHING VESSEL FISHERIES FOR KELP BASS AND BARRED SAND BASS IN THE SOUTHERN CALIFORNIA BIGHT

MILTON S. LOVE and ANDREW BROOKS

Marine Science Institute
University of California
Santa Barbara, California 93106

J. R. RAYMOND ALLY

California Department of Fish and Game
330 Golden Shore, Suite 50
Long Beach, California 90802

We describe the sport fishery for kelp bass, *Paralabrax clathratus*, and barred sand bass, *P. nebulifer*, in the Southern California Bight using catch data from two commercial passenger fishing vessel (CPFV) creel censuses conducted from 1975 to 1978 and 1986 to 1989. In the 1986-1989 census, both species ranked in the top three as measured by catch per unit effort (CPUE). Catch per unit effort of both species was much higher during 1986-1989 than in 1975-1978. The increased catches in the 1980s were likely due to increased abundance, rather than changes in targeting by vessel operators. Sites with highest CPUE for kelp bass contained high relief, both with and without kelp. The highest CPUE occurred at several oil platforms off Santa Barbara. Most of the best fishing sites were located in the northern part of the Southern California Bight, probably because fishing pressure is lower there than to the south. Kelp bass fishing was best in late spring and early fall; catches decreased during the peak of the spawning period. Highest catches of barred sand bass occurred over low relief or sandy bottoms where spawning aggregations occurred. Catch per unit effort was highest during the summer spawning period. Kelp bass were most commonly taken from <30 m. Barred sand bass were most abundant in 10-50 m. Mean lengths of both species taken in the CPFV fishery were essentially unchanged between the 1970s and 1980s; during both periods the majority of the fishes caught were above the legal size limit of 30.5 cm total length.

INTRODUCTION

Kelp bass, *Paralabrax clathratus*, and barred sand bass, *P. nebulifer*, are two very abundant near-shore species in the Southern California Bight. Excluded by law from the California commercial fishery since 1953, these two sport fishes are among the three most important species in the commercial passenger fishing vessel (CPFV)

catch (Ally et al.¹ 1991) and in the top six in private boat landings (Wine² 1979). Commercial passenger fishing vessels may take >1 million kelp bass and >600,000 barred sand bass annually (Ally et al.¹ 1991).

Kelp bass are popular because they strike hard, are relatively easy to hook, are often taken near the surface, and are palatable. Additionally, because kelp bass tend to be found around fouling areas (kelp beds, oil platforms, and high-relief reefs), there is a degree of difficulty landing them, though they are not as difficult to boat as larger species, such as yellowtail, *Seriola lalandi*. Many fishing clubs in southern California charter CPFVs specifically to catch kelp bass. Several CPFV operators note that on slow kelp bass days, many passengers would rather continue fishing for that species than switch to more readily available species, such as the various shallow water rockfishes (*Scorpaenidae*).

Barred sand bass are also popular and sought after by CPFV anglers. Most sand bass are taken during summer, when they aggregate at specific, predictable sites to spawn. During these times, hundreds of fish may be taken per vessel. In general, this bottom-dwelling species requires less skill to hook and land than kelp bass and the fish is held in slightly lower esteem than kelp bass by many anglers. Operators often seek out barred sand bass when their passengers include large numbers of relatively inexperienced fishermen.

Despite considerable previous research, little has been published on the sport fishery for barred sand bass. The kelp bass fishery has not been studied since the 1950s (Young 1963). This paper summarizes a series of studies that fill in gaps in our knowledge.

METHODS

Fishes taken aboard CPFVs were sampled by the senior author and by California Department of Fish and Game (CDFG) personnel from 1975 to 1978 and by CDFG personnel from 1986 to 1989. We randomly sampled regularly scheduled trips by vessels operating from Point Conception to the Mexican border. In the 1970s, 2,234 trips were surveyed and 342,051 fishes identified; in the 1980s, 4,938 trips were sampled and 437,028 fishes identified.

The sampler boarded the vessel at the beginning of the trip and remained aboard to identify; count; and, if possible, measure all fishes caught by passengers. Particular effort was made to count and measure all fishes released. The sampler also noted date, whether the boat fished in Mexican waters, number of anglers (including

¹ Ally, J.R.R., D.S. Ono, R.B. Read, and M. Wallace. 1991. Status of major southern California marine sport fish species with management recommendations, based on analyses of catch and size composition data collected on board commercial passenger fishing vessels from 1985 through 1987. California Department of Fish and Game, Marine Resources Administrative Report No. 90-2.

² Wine, V. 1979. Southern California independent sport fishing survey, annual report no. 3. California Department of Fish and Game, Marine Resources Administrative Report No. 79-3.

crew, if they fished), time fished in each location, and bottom depth.

Each fish was measured to the nearest millimeter. When fishes were brought aboard too rapidly for all to be measured or identified, samplers gave priority to those being returned to the water and measured or identified the remaining fishes at the end of the day.

We estimated relative fish abundances using catch per unit effort (CPUE), defined as number of fish taken per angler-h (where angler-h = number of anglers x number of hours fished). To rank species within each year, we calculated an annual CPUE by dividing total number of fish caught (by species) by total effort during the year. We characterized each fishing site in terms of overall CPUE for kelp bass and barred sand bass by grouping all years in each of the two time periods (1975-1978 and 1986-1989) and dividing the species' total catch at each site by the effort at that site.

During the year (or annually), CPFV operators may shift their emphasis among a variety of species. For instance, while deeper-water rockfishes are targeted during winter, the same vessels may shift to kelp bass in spring and barred sand bass in summer. Also, estimates of seasonal changes in CPUE (and relative abundance) of kelp bass and barred sand bass may be biased by shifts in fishing effort from reefs inhabited by these species to reefs where they are absent. Thus, for our analysis of seasonal changes in CPUE, we included data only from sites that were fished during all seasons and where kelp bass or barred sand bass were taken during at least one season. To compare the CPUE of each species between years, we used only those reefs on which the species occurred during at least 1 of the 8 years. All comparisons were made using standard analysis of variance with individual trips counted as separate observations for calculation of mean CPUEs. Multiple comparison tests were made using the Ryan-Einot-Gabriel-Welsch Multiple F-test (REGWF option, Proc GLM, SAS 1987).

Standard statistical methods were used for other comparisons. We used an approximate t-test, assuming unequal variances, to compare mean lengths between periods. Chi-square was used to compare length-frequency distributions.

RESULTS AND DISCUSSION

Annual Catch Per Unit Effort

Kelp bass were an important recreational species in the 1975-1978 survey, consistently ranking third in CPUE; barred sand bass were less important, ranking between fifth and tenth (Table 1). Catches of both species, but particularly barred sand bass, were relatively low. The maximum CPUE of kelp bass was 0.26; that of barred sand bass, 0.07.

In 1986-1989, kelp bass and barred sand bass were consistently among the top three species in the CPFV catch (Table 2). Kelp bass were particularly important, ranking either first or second in all 4 years. During the same period, barred sand bass ranged from first to third.

Catch rates of both species were higher during 1986-1989 than in 1975-1978

Table 1. Top ten species in the southern California commercial passenger fishing vessel catch from 1975-1978 as measured by the number of fishes taken per angler-h (in parentheses).

1975		1976	
1. Boccacio, <i>Sebastes paucispinis</i>	(0.42)	1. Boccacio	(0.41)
2. Pacific Bonito, <i>Sarda chiliensis</i>	(0.08)	2. Chilipepper	(0.14)
3. Kelp Bass, <i>Paralabrax clathratus</i>	(0.07)	3. Kelp Bass	(0.13)
4. Chilipepper, <i>S. goodei</i>	(0.06)	4. Olive Rockfish	(0.07)
5. Olive Rockfish, <i>S. serranoides</i>	(0.06)	5. Barred Sand Bass	(0.06)
6. Blue Rockfish, <i>S. mystinus</i>	(0.04)	6. Pacific Bonito	(0.06)
7. Chub Mackerel, <i>Scomber japonicus</i>	(0.03)	7. Pacific Barracuda	(0.04)
8. Sablefish, <i>Anoplopoma fimbria</i>	(0.03)	8. Blue Rockfish	(0.03)
9. Pacific Barracuda, <i>Sphyrna argentea</i>	(0.03)	9. Ocean Whitefish, <i>Caulolatilus princeps</i>	(0.02)
10. Barred Sand Bass, <i>Paralabrax nebulifer</i>	(0.03)	10. Vermilion Rockfish, <i>Sebastes miniatus</i>	(0.02)
1977		1978	
1. Chub Mackerel	(0.21)	1. Chub Mackerel	(0.21)
2. Boccacio	(0.20)	2. Boccacio	(0.29)
3. Kelp Bass	(0.17)	3. Kelp Bass	(0.26)
4. Olive Rockfish	(0.13)	4. Chilipepper	(0.13)
5. Chilipepper	(0.11)	5. Pacific Bonito	(0.08)
6. Blue Rockfish	(0.09)	6. Blue Rockfish	(0.06)
7. Barred Sand Bass	(0.07)	7. Olive Rockfish	(0.06)
8. Pacific Bonito	(0.05)	8. Barred Sand Bass	(0.06)
9. Pacific Barracuda	(0.04)	9. White Croaker, <i>Genyonemus lineatus</i>	(0.06)
10. Halfmoon, <i>Medialuna californiensis</i>	(0.04)	10. Pacific Barracuda	(0.03)

(Table 2). This pattern was particularly evident for barred sand bass, where the 1980s CPUE was 5-10 times that of the 1970s. Because kelp bass and barred sand bass are very popular recreational species (actively sought by CPFV operators), and have been for many decades (Frey 1971), we believe low CPUEs in the previous period were due to depressed abundances of these fishes, rather than operators targeting other species.

Kelp bass CPUE differed significantly among years ($F=25.85, P<0.001$). Catch per unit effort was significantly and consistently higher beginning in 1978 (Fig. 1). A similar pattern held for barred sand bass ($F=14.84, P<0.001$) (Fig. 1). Again, CPUE in the 1980s was significantly higher than in the 1970s. It is possible that populations of both kelp bass and barred sand bass benefited from rising ocean temperatures in the mid-1970s and increased recruitment success (Stephens et al. 1994).

Table 2. Top ten species in the southern California commercial passenger fishing vessel catch from 1986-89 as measured by the number of fishes taken per angler-h (in parentheses).

1986		1987	
1. Kelp Bass	(0.44)	1. Chub Mackerel	(0.40)
2. Chub Mackerel	(0.34)	2. Kelp Bass	(0.34)
3. Barred Sand Bass	(0.19)	3. Barred Sand Bass	(0.34)
4. Pacific Bonito	(0.18)	4. Pacific Bonito	(0.21)
5. Pacific Barracuda	(0.13)	5. Pacific Barracuda	(0.16)
6. Boccacio	(0.07)	6. Chilipepper	(0.08)
7. Ocean Whitefish	(0.04)	7. Boccacio	(0.05)
8. White Croaker	(0.04)	8. California Scorpionfish	(0.04)
9. Chilipepper	(0.03)	9. Vermilion Rockfish	(0.04)
10. California Scorpionfish, <i>Scorpaena guttata</i>	(0.03)	10. Ocean Whitefish	(0.03)
1988		1989	
1. Barred Sand Bass	(0.33)	1. Kelp Bass	(0.32)
2. Kelp Bass	(0.28)	2. Chub Mackerel	(0.27)
3. Chub Mackerel	(0.22)	3. Barred Sand Bass	(0.27)
4. Pacific Barracuda	(0.12)	4. Pacific Bonito	(0.16)
5. California Scorpionfish	(0.10)	5. Pacific Barracuda	(0.11)
6. Pacific Bonito	(0.09)	6. California Scorpionfish	(0.10)
7. Chilipepper	(0.06)	7. Boccacio	(0.10)
8. Vermilion Rockfish	(0.06)	8. Chilipepper	(0.08)
9. White Croaker	(0.04)	9. White Croaker	(0.04)
10. Boccacio	(0.04)	10. Vermilion Rockfish	(0.04)

The increased year-class strength of both species in the late 1970s is also reflected in the changes in the proportion of legal-sized fishes (>30.5 cm total length) taken by CPFVs (Fig. 2). The proportion of legal-sized kelp bass was higher in 1975-76 than in 1977-78. The proportion of legal-sized sand barred bass remained high through 1977, then decreased slightly in 1978. We believe these shifts in size composition of the catch were due to increased numbers of small bass throughout the Southern California Bight (Stephens et al. 1994), reflecting substantial recruitment of kelp bass in the late 1970s, and the lesser success of barred sand bass during the same period, rather than a decrease in the number of larger fish due to fishing.

Geographic Distribution

Several productive kelp bass fishing sites exist in the Southern California Bight. Kelp bass tend to be found over high-relief, often kelp-covered, substrates. Areas off rocky headlands, such as Palos Verdes and Point Dume, seem particularly favorable. Excluding sites with low (<100 angler-h) effort, 22 sites had CPUEs >1.0 kelp bass per angler-h (Fig. 3, Table 3). While all these sites contained high relief, some were natural formations and some were artificial. Kelp was present on most, but not all, of the most important sites.




Figure 1. Annual mean catch per angler-h of kelp bass and barred sand bass taken in the southern California commercial passenger fishing vessel fishery, 1975-1978 and 1986-1989. Horizontal lines below the abscissa indicate statistically significant groupings using the Ryan-Einot-Gabriel-Welsch Multiple F-test. Error bars are \pm one standard error about the mean.

With the exception of one site (Cardiff-By-The-Sea, about 40 km north of San Diego), all of the most productive kelp bass sites were either along the mainland in the northernmost part of the Southern California Bight or around the Channel Islands. Of the 22 sites, 11 were mainland ones within 30 km of Santa Barbara. Including the northern Channel Island locations with the mainland sites, 17 of 22 sites are in this region. The highest CPUE of any of these areas was 2.4 kelp bass per




Figure 2. The proportion of legal-sized (>30.5 cm) kelp bass and barred sand bass taken in the southern California commercial passenger fishing vessel sport fishery, 1975-1978 and 1986-1989. Error bars represent 95% confidence limits calculated as in Rohlf and Sokal (1981).

angler-h at the adjoining oil platforms Hilda and Hazel (data from these platforms were pooled) situated in about 30 m of water, 3.5 km offshore of Summerland, 10 km east of Santa Barbara. Kelp does not grow on these platforms, nor adjacent to them. In contrast, the next highest CPUE, 2.3 kelp bass per angler-h, was obtained at Naples Reef, a high-relief natural formation (usually kelp covered) located at a depth of 8-12 m, 1.6 km offshore of Naples, 24 km west of Santa Barbara.

This pattern in CPUE suggests that fishing success is inversely related to fishing effort. The northern part of the bight (from Ventura north) is lightly fished when compared to areas from Los Angeles south (Fig. 4). This southern area is more




Figure 3. Catch per angler-h (CPUE) for kelp bass taken in the Southern California Bight by the commercial passenger fishing vessel fishery, 1986-1989. Due to space limitations, the following Santa Cruz Island locations with CPUE >1.0 are illustrated as one site: Chinese Harbor, Smugglers Cove, San Pedro Point, Scorpion Arch, and Yellow Banks. One other high CPUE site, Avalon Bay, Santa Catalina Island, is not shown.

heavily populated and contains larger recreational fishing fleets. Moreover, the southern areas are less affected by storms and have more fishable days. The same reasoning helps explain the productivity of most of the San Clemente Island sites. That island is one of the most remote in southern California and, except for the leeward side, receives less attention from recreational fishermen than sites closer to ports.

Avalon Bay on Santa Catalina Island ranked third in CPUE, even though Santa Catalina Island is heavily fished. Avalon, the only developed harbor on the island, is fished continuously by both CPFVs and private vessels. One possible reason for the abundance of kelp bass in and around the bay is the heavy daily feeding they receive from local glass-bottom tour boats. As these vessels cruise the local reefs, they toss food into the water, which brings kelp bass under the vessel, where they are visible to passengers. Feeding may occur hourly during daytime. The large numbers of kelp bass in the bay may be attracted to, and held in the vicinity, by this continual feeding. Kelp bass larval recruitment is higher at Catalina Island than at a mainland site (Carr 1994) and this also may help explain the phenomenon. In addition, there is a marine refuge close to Avalon, where no fishing is allowed. Kelp bass from the refuge may wander over to Avalon and enter the recreational fishery.

In contrast to kelp bass, the areas with highest CPUE for barred sand bass were distributed along the mainland from Ventura into Baja California (Fig. 5, Table 4). Nine sites yielded CPUEs >1.0 barred sand bass per angler-h. Barred sand bass prefer lower relief sites away from rocky headlands; they are often found in mid-bay (e.g., Santa Monica Bay) or along sandy stretches of coast. Most fishing effort for barred

Table 3. Commercial passenger fishing vessel kelp bass fishing sites in the Southern California Bight with catch per angler-h ≥ 1.0 . Based on 1986-89 data, excluding sites with <100 angler-h effort. Except Cardiff-by-the-Sea (located near Oceanside), all mainland sites are located within 30 km of Santa Barbara.

<u>Rank</u>	<u>Site name</u>	<u>Effort</u>	<u>CPUE</u>
1	Oil Platforms Hilda and Hazel	529.8	2.4
2	Naples Reef	298.2	2.3
3	Avalon Bay, Santa Catalina I	210.2	2.2
4	Chinese Harbor, Santa Cruz I	222.7	1.7
5	Smugglers Cove, Santa Cruz I	707.5	1.7
6	East End, San Clemente I	146.4	1.7
7	East End, Santa Cruz I	142.7	1.6
8	West End, San Clemente I	328.2	1.5
9	Camby's Reef	441.7	1.5
10	Mohawk Reef	169.0	1.5
11	Horseshoe Kelp ^a	550.2	1.4
12	Ellwood	240.9	1.4
13	Hendry's Beach	130.7	1.2
14	San Pedro Pt., Santa Cruz I	878.9	1.1
15	Scorpion Arch, Santa Cruz I	539.1	1.1
16	Isla Vista Reef	208.4	1.1
17	Rincon Oil Platforms	210.2	1.1
18	Horseshoe Reef	673.6	1.1
19	Yellowbanks, Santa Cruz I	1,579.0	1.1
20	Cardiff-By-The-Sea	555.3	1.0
21	Leeward Side, San Clemente I	1,035.3	1.0
22	Flag Reef	634.4	1.0

^aIn southern California, there are two sites called "Horseshoe Kelp". This one is a small reef located just south of Santa Barbara, not the large structure situated offshore of the San Pedro-Long Beach Harbor.

sand bass occurs during the late spring-early fall spawning season, when CPFV operators target large spawning aggregations. During the non-reproductive period, the fish tend to be found in small groups near low relief. Thus, sites with the highest CPUE may represent spawning areas for the species. In general, these are areas of low relief or hard bottom, often in 20-40 m. While barred sand bass are abundant all along the coast, they are relatively rarely caught around the islands of the bight, despite a great deal of fishing pressure and a fair amount of suitable habitat, at least around Santa Catalina and Santa Cruz islands.

Depth Distribution

While kelp bass were taken at sites with bottom depths from <10 to >50 m, they were most abundant at depths <30 m (Fig. 6). Catches decreased in 31-40 m and few fish were taken in deeper water. Seasonal catch patterns implied an offshore




Figure 4. Mean total effort expended over the 4-year period, 1985-1988, in seven regions of the Southern California Bight. Error bars represent \pm one standard error about the mean. Numbers in parenthesis refer to the number of locations fished in each region.

movement of fish during spring, at least from the shallowest depths. Of all depths and seasons, CPUE's were highest in <10 m during winter. By spring, catches in these depths had decreased by almost a factor of 3, corresponding with movements to the deeper water noted by Quast (1968). This offshore movement may be related to this species' small-scale spawning migrations (Feder et al. 1974). Since our analysis was only based on sites harboring kelp bass, the declining CPUE in 0-10 m during spring and summer is not due to a shift in CPFV fishing sites.

Barred sand bass were taken in depths ranging from <10 to >50 m (Fig. 6), with highest catches in 11-50 m. Depths of the summer spawning aggregations (centered between 21 and 40 m) are apparent. Compared to other seasons, catches were considerably higher during the summer months in 21-40 m. Some slight winter inshore movement may occur, as catches during the winter in the shallowest areas (<10 m) were higher than in other seasons.



Figure 5. Catch per angler-h (CPUE) for barred sand bass taken in the Southern California Bight by the commercial passenger fishing vessel fishery, 1986-1989.

Table 4. Commercial passenger fishing vessel barred sand bass fishing sites in the Southern California Bight with catch per angler-h ≥ 1.0 . Based on 1986-89 data, excluding sites with < 100 angler-h effort. Sites are designated by the closest large city.

Rank	Site name	Effort	CPUE
1	Rincon Pt. - Loon Pt. (Ventura)	147.0	4.5
2	Silver Strand (San Diego)	508.0	1.6
3	"Sand Bass Junction" (Long Beach)	1,443.4	1.4
4	San Clemente Reef (San Clemente)	1,328.0	1.4
5	Ventura Harbor-Pitas Pt. (Ventura)	3,434.5	1.3
6	Seal Beach (Long Beach)	772.3	1.3
7	Shale Spot (Ventura)	1,225.0	1.3
8	Huntington Flats (Huntington Beach)	12,117.0	1.1
9	Inner Santa Monica Bay (Santa Monica)	3,088.0	1.1

Total length for both kelp bass and sand bass exhibited a positive relationship with depth of capture during the 1970s (kelp bass: slope = 0.29, $F = 16.76$, $P < 0.001$; barred sand bass: slope = 3.96, $F = 631.60$, $P < 0.001$) and the 1980s (kelp bass: slope = 0.31, $F = 92.74$, $P < 0.001$; barred sand bass: slope = 0.74, $F = 381.89$, $P < 0.001$), suggesting that small individuals are caught in shallower depths. In spite of the significance of these regressions, depth actually explained very little of the total variation in length of the catch of either species during either the 1970s (kelp bass: $r^2 = 0.001$; barred sand bass: $r^2 = 0.059$) or the 1980s (kelp bass: $r^2 = 0.003$; barred sand bass: $r^2 = 0.017$) and fish of all sizes were taken in all depths sampled.

Size Frequencies

Mean total length of kelp bass taken in the CPFV fishery during the years 1975-1978 was 33.6 cm. This value increased slightly to 33.7 cm during the period 1986-1989 (Fig. 7), but this increase was not statistically significant ($t = 1.55$, $P > 0.05$). Mean total length of barred sand bass decreased significantly from 35.7 cm in the 1970s to 35.3 cm during the 1980s ($t = 4.47$, $P < 0.001$) (Fig. 8). Such a small decrease in size is probably not biologically significant.

To determine if kelp bass and barred sand bass were subject to growth overfishing (manifested as a decrease in the proportion of large fish in the catch), we tested for differences between the 1970s and 1980s in the percentage of the catch that was >30.5 cm (the minimum legal length). Length-frequency distributions of the catches




Figure 6. Variability in catch/angler-h with depth for kelp bass and barred sand bass in the southern California commercial passenger fishing vessel fishery, 1986-1989. Error bars are \pm one standard error about the mean.




Figure 7. Length frequency distribution of kelp bass caught in the southern California commercial passenger fishing vessel fishery, 1975-1978 and 1986-1989. Vertical dashed lines represent the minimum legal size limit and the black circle indicates the sample mean length, \pm one standard deviation about the mean.

(Fig. 7 and 8) indicate that the fishery did not crop out the larger fish between the 1970s and 1980s. In fact, the percentage of the catch above legal size increased significantly, from 65% to 77% for kelp bass ($\chi^2 = 1,532.0$, $P < 0.0001$) and from 83% to 88% for barred sand bass ($\chi^2 = 211.0$, $P < 0.001$).

The proportion of legal-sized to sublegal fish likely reflects the success of recruitment in previous years. For instance, the sharp decrease in the proportion of legal-sized kelp bass between 1976 and 1978 closely parallels increased recruitment success beginning in 1976 off Redondo Beach (Stephens et al. 1986).




Figure 8. Length frequency distribution of barred sand bass caught in the southern California commercial passenger fishing vessel fishery, 1975-1978 and 1986-1989. Vertical dashed lines represent the minimum legal size limit and the black circle indicates the sample mean length, \pm one standard deviation about the mean.

Seasonality

During 1986-89, kelp bass CPUE was highest in late spring and fall (Fig. 9). Catches were low from January through March, rose to a May-June peak, then declined during summer to an August low. A second, and higher, peak occurred in October. Catches during the October peak were 4 times those of the February low.




Figure 9. Monthly variability in mean catch/angler-h of kelp bass and barred sand bass taken in the southern California commercial passenger fishing vessel sport fishery during the years 1986-1989. Error bars are \pm one standard error about the mean.

The low catches during winter may be due to the effects of lower water temperatures on metabolism and feeding. Other factors may be responsible for the spring and fall peaks and summer low. Quast (1968) reported a similar pattern in CPUE of kelp bass taken from June-October 1958 aboard a single CPFV fishing off San Diego. He found no significant correlations between CPUE and environmental factors (including tides, weather, lunar phases, and surface temperatures) and suggested that kelp bass

spawning condition might have been a significant factor. He theorized that kelp bass feed more heavily during spawning. However, based on his data, most spawning appears to occur from July-September and the spring catch maximum comes just before, and the October maximum just after, the peak spawning period. Rather than feeding more during spawning season, kelp bass may feed less.

Barred sand bass catches showed a different pattern. Catches were relatively low throughout the year, except for a summer (July-August) peak (Fig. 9). Catch per unit effort varied by a factor of 4 between the May low and July high. As mentioned before, this species is a summer spawner and the catch pattern reflects the targeting of spawning aggregations with their greatly increased concentrations of barred sand bass.

Future of the Fishery

The data presented here demonstrates the importance of kelp bass and barred sand bass to the CPFV fleet. Even during the 1970s, when catches of both species were considerably lower than during the 1980s, these species ranked high in CPUE relative to other fishes. The CPFV fishery frequently targets both kelp bass and barred sand bass, while other species represent incidental catch.

Kelp and barred sand bass fisheries appear to be in good condition. The bass populations (as indexed by CPUE), bolstered by minimum length regulations, increased between the 1970s and 1980s, while the overall size structure of the catch has remained relatively stable. Thus, the bass fisheries are currently managed to optimize the quantity of bass caught, rather than size.

However, some anglers feel that there are not enough large, "trophy" fish in the population and that actions to improve the quality (as opposed to the quantity) of the bass fishery, particularly for kelp bass, are necessary. Their solution is to return virtually all bass caught. This is an almost revolutionary concept for the CPFV anglers of southern California. Traditionally, these anglers have kept virtually all the legal-sized popular species caught.

Our observations over the last 20 years show that there were several reasons for this "catch and keep" phenomenon. First, for many anglers, the fish they catch are a food source. Second, there is a competitive aspect to catching the most fish; full sacks mark the successful angler. In addition, CPFV operators and crews encourage fish retention in order to (i) bolster the number of fishes taken and make all passengers happy, (ii) attract future customers when "fish counts" of each landing are listed in the daily newspapers, and (iii) maximize income that comes from cleaning fish at the end of the trip.

Opposing these long-held behaviors are a small, but growing, group which points out that returning fish is no longer an unusual circumstance among other groups of recreational fishermen. We found that several CPFV operators are encouraging their passengers to release unwanted kelp bass (M. McCrae, F/V *SEAHAWK*, Sea Landing, Santa Barbara, California, pers. comm.). Mr. McCrae explains over the vessel's public address system that the reef they are fishing is small; can be easily fished out; and that, if they release the kelp bass, the fish may be there the next time they fish that reef.

Kelp bass is an excellent species for a catch and release program. It is taken in shallow water, so the inflated swimbladder problems found in deeper-dwelling species (such as rockfishes) are nonexistent. It is also very hardy; our observations show that almost all kelp bass hooked can be released without fatally injuring the fish. Finally, at least some kelp bass populations seem to be resident, so a catch and release program should show results within a relatively few years.

ACKNOWLEDGMENTS

The research was supported by the Federal Aid for Sport Fish Restoration Program and the National Biological Service, contract number 1445-CA-0995-0386. We thank R. Collins for providing the 1970s creel census data. L. Allen and C. Valle critically reviewed drafts of the manuscript. We also thank two anonymous reviewers for very constructive comments.

LITERATURE CITED

- Carr, M.H. 1994. Effects of macroalgal dynamics on recruitment of a temperate reef fish. *Ecology* 75:1320-1333.
- Feder, H.M., C.H. Turner, and C. Limbaugh. 1974. Observations on fishes associated with kelp beds in southern California. California Department of Fish and Game, Fish Bulletin 160.
- Frey, H. (editor). 1971. California's living marine resources and their utilization. California Department of Fish and Game, Sacramento, California, USA.
- Quast, J.C. 1968. Observations on the food and biology of the kelp bass, *Paralabrax clathratus*, with notes on its sportfishery at San Diego, California. Pages 81-108 in: W.J. North and C.L. Hubbs, editors. Utilization of kelp-bed resources in southern California. California Department of Fish and Game, Fish Bulletin 139.
- Rohlf, F.J. and R.R. Sokal. 1981. Statistical tables, 2nd edition. D.W. Freeman and Co., New York, New York, USA.
- SAS. 1987. SAS procedures guide, release 6.04 edition. SAS Institute Inc., Cary, North Carolina, USA.
- Stephens, J.S., Jr., G.A. Jordan, P.A. Morris, M.M. Singer, and G.E. McGowen. 1986. Can we relate larval fish abundance to recruitment or population stability? A preliminary analysis of recruitment to a temperate reef. California Coastal Oceanic Fisheries Investigations Reports 27:65-83.
- Stephens, J.S. Jr., P. Morris, D. Pondella, T. Koonce, and G. Jordan. 1994. Overview of the dynamics of an urban artificial reef fish assemblage at King Harbor, California, USA, 1974-1991: A recruitment driven system. *Bulletin of Marine Science* 55:1224-1239.
- Young, P.H. 1963. The kelp bass (*Paralabrax clathratus*) and its fishery, 1947-1958. California Department of Fish and Game, Fish Bulletin 122.

Received: 3 December 1994

Accepted: 1 November 1995

CONDITION AND REPRODUCTIVE PERFORMANCE OF FEMALE MULE DEER IN THE CENTRAL SIERRA NEVADA

T. J. TAYLOR

P.O. Box 191

June Lake, California 93529

I determined reproductive performance in relation to age and physical condition of 58 female Rocky Mountain mule deer, *Odocoileus hemionus hemionus*, collected from the West Walker winter range in northern Mono County, California, and southwestern Douglas County, Nevada during March 1993 and March 1994. Among adult females, pregnancy and fetal rates were 86% and 1.52 fetuses/female in 1993 and 88% and 1.56 fetuses/female in 1994. Bled carcass weights, eviscerated carcass weights, and kidney fat indices of adult females were greater in 1994, indicating that deer were in better condition than in 1993. Litter category was not related to age or female body condition. The overall sex ratio of fetuses was 107 males:100 females. Reproductive potential in the West Walker deer herd was comparable to that reported for other nutritionally stressed mule deer populations. My study suggested that low productivity was in response to drought-induced changes in habitat quality.

INTRODUCTION

During 1987-1994, Rocky Mountain mule deer, *Odocoileus hemionus hemionus*, populations in the eastern Sierra Nevada, California, experienced poor fawn survival and low recruitment. For example, fall fawn ratios for the West Walker (WW) deer herd in northern Mono County averaged 26 fawns:100 females, while spring ratios averaged 20 fawns:100 females (R. Thomas, California Department of Fish and Game, pers. comm.). This low recruitment was concomitant with 7 consecutive years of drought, which subjected this deer population to repeated episodes of nutritional stress.

Nutrition is an important factor influencing productivity of deer populations (Verme 1967, McCullough 1979) because it affects the proportion of females that become pregnant, as well as ovulation rates (Sadleir 1987, Folk and Klimstra 1991). These effects are well documented for free-ranging (Morton and Cheatum 1946, Julander et al. 1961, Kucera¹ 1988) and captive deer (Verme 1965, 1967; Ozoga and Verme 1982). Research consistently has shown that deer on good quality ranges have higher rates of ovulation, conception, and pregnancy than deer on poor ranges.

An understanding of the relationships between body condition and reproductive performance of wild ungulate populations is necessary for their management

¹ Kucera, T.E. 1988. Ecology and population dynamics of mule deer in the eastern Sierra Nevada, California. Ph.D. Dissertation, University of California, Berkeley, California, USA.

(Saltz et al. 1992). Therefore, a high priority in any deer herd where fawn production is suboptimal should be to evaluate the physical condition of females during breeding and pregnancy (Connolly 1981, Saltz et al. 1992).

Body fat is the component most often associated with animal condition and can be used to index animal response to nutritional and climatic stressors (Robbins et al. 1974, Torbit et al. 1985). Various fat indices have been developed to estimate body condition, including bone marrow fat (Cheatum 1949; Riney 1955; Ransom 1965, 1967), kidney fat (Riney 1955; Ransom 1965, 1967; Batcheler and Clark 1970; Van Vuren and Coblentz 1985), and visual scoring methods (Ransom 1965, Kistner² 1976). I used kidney fat indices and reproductive tracts to determine the physical condition and reproductive potential of adult female mule deer collected from the WW deer herd during March 1993 and March 1994. My objectives were to (i) assess the effects of a prolonged drought, followed by an unusually severe winter, on the spring condition of female mule deer and (ii) compare the effects of an unusually severe and an unusually mild winter on deer condition and productivity. This descriptive study is intended to provide information on mule deer condition and reproduction in order to facilitate a better understanding of interactions that occur between the WW deer herd and its environment.

STUDY AREA AND METHODS

During 18-19 March 1993 and 15-16 March 1994, free-ranging female mule deer were collected from WW winter ranges in northern Mono County, California, and southwestern Douglas County, Nevada (Fig. 1). Winter range of the WW herd encompasses approximately 780 km² at elevations from 1,530 to 2,550 m. Dominant plant communities (following Mayer and Laudenslayer 1988) on the winter range include bitterbrush, *Purshia* sp.; sagebrush, *Artemisia* sp.; and pinyon pine, *Pinus monophylla*. Descriptions of WW herd ecology, winter range vegetation, climate, and topography have been reported by Thomas³ (1985), Loft et al. (1989), and Taylor⁴ (1993).

Deer were collected by two-person teams that shot the first identifiable adult or yearling female in each group of deer encountered, regardless of the animal's apparent body condition. Animals were shot in the head, neck, or thorax with a high-powered rifle. Carcasses were transported to a field processing station where they were weighed to the nearest kilogram (bled carcass weight, BCW) using a spring scale. External body measurements (chest girth, left hind foot length, and contour length) to the nearest centimeter were recorded. Animals were eviscerated and reproductive tracts (uterus and ovaries), right kidneys, right femurs, and lower jaws

²Kistner, T.P. 1976. Evaluating physical condition of deer. Oregon Department of Fish and Wildlife, Portland, Oregon, USA.

³Thomas, R.D. 1985. Management plan for the West Walker deer herd. California Department Fish and Game, Bishop, California, USA.

⁴Taylor, T.J. 1993. West Walker deer herd progress report 2. California Department Fish and Game, Bishop, California, USA.



Figure 1. Locations of deer collection areas on the West Walker deer herd winter range in Mono County, California and Douglas County, Nevada, 1993-1994.

were extracted. Ages were estimated by tooth wear and replacement (Larson and Taber 1980). After field necropsies were completed, deer were weighed to the nearest kilogram to determine eviscerated carcass weights (ECW). The kidney fat index (KFI, Riney 1955) was calculated by dividing the fresh weight of kidney fat by the fresh weight of the fat-free kidney, multiplied by 100.

Ovaries were sectioned at 5-mm intervals and examined macroscopically for corpora lutea of pregnancy (CLP), which were used to estimate ovulation rates. I used a fetus scale (Forestry Suppliers Inc., Jackson, Mississippi, USA) to determine forehead-rump lengths of fetuses in order to estimate fetal age (Hudson and

Browman 1959) and conception and fawning dates. Conception date was back-calculated from estimated fetal age. Approximate date of parturition was determined by adding 204 d (Anderson 1981) to conception date. Weather data were obtained from the National Oceanic and Atmospheric Administration weather station in Coleville, California.

I used standard techniques for statistical testing with $\alpha = 0.05$. Analysis of variance was used to test for differences in mean reproductive characteristics and condition indices between years and among collection areas and to examine the relationship between "litter category" (number and sex composition of litters, i.e., single female, twin females, single male, etc.) and BCW, ECW, and KFI. Kidney fat indices were normalized by logarithmic transformation prior to analysis. I tested for deviations from the expected 1:1 sex ratio among fetuses with a binomial test (Siegel 1956); deviations from the expected distribution of sex ratios among litter categories were tested using chi-square analyses. I also used chi-square tests to examine differences in fetal sex ratio between years and among collection areas. Means and standard errors were calculated from untransformed data.

RESULTS

Deer (29 each year) were collected from three primary areas within the range that supported deer concentrations during all winters. A total of 20 deer (10 each year) were collected from Little Antelope Valley (LAV), California; 20 deer (10 each year) were collected from the east side (ES) of Antelope Valley near the base of the Wellington Hills, Nevada; and 18 deer (nine each year) were collected from the vicinity of Topaz Lake (TL), Nevada (Fig. 1).

Winter weather in 1992-93 was severe, with 180 cm of snow and average minimum temperatures of -16°C in January. In comparison, the winter of 1993-94 was mild, with 81 cm of snow and minimum temperatures averaging -6°C in January.

Pregnancy and fetal rates were similar in 1993 and 1994 and mean age (4.9 yr in 1993 and 5.2 yr in 1994) of adult females did not differ significantly between years ($F = 0.227$; 1, 54 df; $P = 0.972$). Among adult females examined, 25 of 29 (86%) were pregnant in 1993 and 24 of 27 (88%) were pregnant in 1994 (Table 1). Mean fetal rates of 1.52 fetuses per adult female in 1993 and 1.56 fetuses in 1994 did not differ between years ($F = 0.357$; 1, 54 df; $P = 0.553$), and mean fetal rates did not differ among collection areas ($F = 0.001$; 2, 53 df; $P = 0.999$) (Table 1). Of the 29 adult females collected in 1993 for which ovaries were examined, 46 CLP resulted in 44 viable, implanted fetuses. Of the 27 adults collected in 1994, 45 CLP resulted in 43 viable, implanted fetuses.

Estimated breeding and parturition dates were similar in both years. Ages of fetuses in 1993 indicate that breeding occurred between 23 November and 30 December, with a median date of 6 December. In 1994, breeding occurred between 28 November and 14 December, with a median date 7 December. Predicted parturition for deer examined in 1993 ranged from mid-June to mid-July; the median was 2 July. In 1994, predicted parturition ranged from 20 June to 7 July; the median was 28 June.

Table 1. Frequency of fetuses in adult Rocky Mountain mule deer collected from the West Walker deer herd winter range in March 1993 and March 1994, Mono County, California and Douglas County, Nevada. LAV = Little Antelope Valley, ES = East Side, TL = Topaz Lake.

	<u>No. of does with litter size of</u>				<u>Total</u>	<u>Total</u>	<u>Mean</u>	<u>Mean.</u>
<u>Location</u>	<u>0</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>no. of</u>	<u>no. of</u>	<u>fetuses</u>	<u>fetuses/</u>
					<u>fetuses</u>	<u>does</u>	<u>per doe</u>	<u>pregnant</u>
								<u>doe</u>
<u>1993</u>								
LAV	0	4	6	0	16	10	1.60 ± 0.52	1.60 ± 0.52
ES	3	1	5	1	14	10	1.40 ± 1.07	2.00 ± 0.58
TL	1	2	6	0	14	9	1.55 ± 0.72	1.75 ± 0.46
<u>1994</u>								
LAV	0	2	7	0	16	9	1.77 ± 0.44	1.77 ± 0.44
ES	2	2	5	0	12	9	1.33 ± 0.86	1.71 ± 0.49
TL	1	1	7	0	15	9	1.67 ± 0.79	1.88 ± 0.35

Deer were in better condition in 1994 than in 1993 (Table 2). Mean BCW of adult females (44.5 kg in 1993 and 50.9 kg in 1994) differed between years ($F = 13.79$; 1, 54 df; $P < 0.001$); however, mean BCW did not differ among collection areas ($F = 0.170$; 2, 53 df; $P = 0.193$), and year by area interaction was not significant ($F = 0.360$; 1, 54 df; $P = 0.700$). Eviscerated carcass weights of adult females followed a similar trend, differing between years ($F = 25.57$; 1, 54 df; $P < 0.001$), but not among collection areas ($F = 2.05$; 2, 53 df; $P = 0.141$); year by area interaction also was not significant ($F = 0.095$; 1, 54 df; $P = 0.910$). Mean KFI was significantly lower in 1993 (9.3%) than in 1994 (36.0%) ($F = 29.53$; 1, 54 df; $P < 0.001$), but did not differ among the three collection areas ($F = 0.939$; 2, 53 df; $P = 0.399$); year by area interaction was not significant ($F = 0.124$; 1, 54 df; $P = 0.884$).

West Walker females produced more male than female fetuses. However, the overall sex ratio of fetuses, 1.07:1.00 in favor of males (45 males:42 females), was not significantly different from unity ($Z = -0.21$, $P = 0.83$) (Table 3). Single fetuses were present in 12 of 49 pregnant females, 36 carried twins, and one had triplets (Table 1). Among the 12 litters of singletons, four fetuses were male and eight were female. Among the 36 sets of twins, 18 were of mixed sex, 9 were twin females, and 9 were twin males. These frequencies of sex ratio categories were not different from random expectation ($\chi^2 = 4.5$, 2 df, $P = 0.105$). Sex ratios of fetuses in 1993 and 1994 did not differ ($\chi^2 = 0.57$, 1 df, $P = 0.451$). Overall, sex ratio categories did not differ among collection areas ($\chi^2 = 5.28$, 2 df, $P = 0.072$), but in 1994 LAV and TL females produced significantly more male fetuses than did ES females ($\chi^2 = 11.3$, 2 df, $P = 0.004$) (Table 3). I found no relationship between litter category and maternal age ($F = 1.55$; 5, 43 df; $P = 0.208$), nor between litter category and BCW ($F = 1.61$; 5, 43 df; $P = 0.178$), ECW ($F = 0.846$; 5, 43 df; $P = 0.52$) and KFI ($F = 1.63$; 5, 43 df; $P = 0.173$).

Table 2. Mean bled carcass weight (BCW), eviscerated carcass weight (ECW), and kidney fat index (KFI) for 56 adult female Rocky Mountain mule deer collected on the West Walker deer herd winter range in March 1993 and March 1994, Mono County, California and Douglas County, Nevada. LAV = Little Antelope Valley, ES = East Side, TL = Topaz Lake.

<u>Location</u>	<u>n</u>	<u>BCW (kg)</u>		<u>ECW (kg)</u>		<u>KFI (%)</u>	
		<u>\bar{X}</u>	<u>SD</u>	<u>\bar{X}</u>	<u>SD</u>	<u>\bar{X}</u>	<u>SD</u>
<u>1993</u>							
LAV	10	44.1	4.52	31.0	3.28	7.2	3.05
ES	10	42.6	5.07	30.2	2.64	9.2	5.88
TL	9	47.1	7.57	34.1	5.04	11.4	9.09
Total	29	44.5	5.90	31.7	3.97	9.3	6.38
<u>1994</u>							
LAV	9	49.7	3.82	35.7	2.24	45.4	27.22
ES	9	51.0	6.16	36.8	4.47	32.2	24.05
TL	9	52.1	4.39	38.6	4.08	30.4	10.72
Total	27	50.9	4.75	37.0	3.79	36.0	22.09

Table 3. Sex of fetuses from 49 adult female Rocky Mountain mule deer collected on the West Walker deer herd winter range in March 1993 and March 1994, Mono County, California and Douglas County, Nevada. LAV = Little Antelope Valley, ES = East Side, TL = Topaz Lake.

<u>Location</u>	<u>Male (N)</u>	<u>Female (N)</u>	<u>% Male</u>
<u>1993</u>			
LAV	7	9	43.7
ES	7	7	50.0
TL	7	7	50.0
Total	21	23	47.7
<u>1994</u>			
LAV	10	6	62.5
ES	2	10	16.6
TL	12	3	80.0
Total	24	19	55.8

DISCUSSION

Reproductive rates observed during this study were similar to those reported by Bischoff (1958), Jordan⁵ (1967), and Kucera¹ (1988) for other central Sierra Nevada mule deer populations. I detected no significant differences in measures of reproduction between years, despite significant increases in measures of body condition in 1994. Fetal rates during this study were 1.52 fetuses/doe in 1993 and 1.56 fetuses/doe in 1994; mean KFI's increased from 9% in 1993 to 36% in 1994.

⁵ Jordan, P.A. 1967. Ecology of migratory deer in the San Joaquin River drainage. Ph.D. Dissertation, University of California, Berkeley, California, USA.

In the eastern Sierra Nevada, Taylor⁶ (1991) documented reproductive rates on two mule deer winter ranges of 1.88 and 1.93 fetuses/doe when KFI's averaged 32% and 25%, respectively. Kucera¹ (1988) found low reproductive rates (1.06-1.42 fetuses/female) among mule deer when mean KFI's averaged 10-27%; a higher reproductive rate (1.88 fetuses/female) was observed with mean KFI >60%. Taylor⁷ (1988) reported that mule deer from the CasaDiablo herd had reproductive rates of 1.74 and 1.70 fetuses/doe when KFI's averaged 41% and 37%, respectively. These incongruous data suggest that KFI may not be an accurate basis for determining the effects of nutritional status on deer productivity when comparing disparate populations.

I surmise that increases in deer condition during 1994 were largely due to the wet winter of 1992-93, which enhanced forage production during the spring and summer of 1993, and increased forage availability during the winter of 1993-94. Kucera¹ (1988) reported large increases in forage production on two mule deer winter ranges in the eastern Sierra Nevada following winters of heavy precipitation, which in turn were mirrored by high measures of condition and reproduction. In contrast, my findings indicate that deer productivity in 1994 was not measurably affected by significant increases in body condition. Following the winter of 1992-93, deer may have so exhausted their body reserves that they were not able to attain the threshold of body condition necessary to enhance their reproductive potential, despite greater forage availability during spring and summer 1993. In years of high forage availability that follow extreme winters, it may not be possible for some deer to achieve enhanced reproductive performance.

The cumulative effects of drought on summer range forage production presumably had a major influence on deer reproduction in 1993. Of 29 adult females, three were without corpora lutea and seven had one corpus luteum each for an ovulation rate of 1.59 CLP per female. This low ovulation rate indicated that summer ranges occupied by WW deer were of poor nutritional quality and, presumably, were inadequate to allow female deer to achieve peak body condition prior to the breeding season. Short (1981) surmised that when summer ranges provide low quality forage because of drought or other factors, the high metabolic requirements of females are not met and requirements for reproduction are not satisfied. Bertram⁸ (1984) reported a pronounced decline in reproductive potential among female mule deer collected from the North Kings herd following the dry spring, summer, and fall of 1977. Several other researchers (Robinette et al. 1955, Swank 1958, Taber and Dasmann 1958, Julander et al. 1961) determined that summer forage nutrition influenced ovulation rates and the number of fetuses produced per female.

⁶Taylor, T.J. 1991. Ecology and productivity of two interstate deer herds in the eastern Sierra Nevada: East Walker-Mono Lake deer herd study. California Department of Fish and Game, Bishop, California, USA.

⁷Taylor, T.J. 1988. The Casa Diablo deer herd: Reproduction and condition 1987-1988. Casa Diablo deer herd study. California Department of Fish and Game, Bishop, California, USA.

⁸Bertram, R.C. 1984. The North Kings deer herd study. California Department of Fish and Game, Fresno, California, USA.

Mean KFI of 9% observed in March 1993 is similar to values reported by Kucera¹ (1988) and Anderson et al. (1972) for other nutritionally stressed mule deer populations. KFI values <15% indicate essentially no fat, and represent the point when animals begin to mobilize femur marrow fat for energy (Ransom 1965, Pojar and Reed 1974, Kie et al. 1983). Animal condition, as indexed by KFI and BCW, was lowest following the severe winter of 1992-93, when snow accumulations on primary winter ranges (1,700 m elevation) totaled 180 cm and lasted from 7 December to 3 March. This prolonged snow cover buried sources of forage and, when coupled with persistent low temperatures, resulted in widespread starvation (T.J. Taylor, unpubl. data) and decreased maternal nutrition and productivity among surviving females. Severe winters have been associated with declines in deer condition and productivity (Leach 1956, Gilbert et al. 1970, Gill 1972, Hall⁹ 1973, Wallmo et al. 1977). During a severe winter, net productivity is influenced not only by deer lost to starvation, but also by the poor condition of surviving females, a situation that results in high postnatal fawn losses (Robinette 1976).

The preponderance of males produced by ES and TL deer sampled in 1994 may be the result of small sample sizes that can result in unusual sex ratios (Thomas et al. 1989). Although Robinette et al. (1957) found that mule deer on poor range produced more male fawns, my results do not suggest a male-biased sex ratio in the nutritionally stressed WW mule deer herd.

Reproductive potential in the WW deer herd during 1993 and 1994 was comparable to that reported for other nutritionally stressed mule deer populations throughout the western United States (Anderson et al. 1972, Kucera¹ 1988). This low productivity was likely in response to drought-induced changes in habitat quality, which was compounded by severe winter conditions in 1992-93. During periods of drought, the first step managers should take to maintain deer herd productivity is to increase the quality of the food supply. Therefore, I recommend management practices on WW herd winter ranges and holding areas that promote and ensure access to late season growth of succulent forage on irrigated pasture. Garrott et al. (1987) found that mule deer in northwest Colorado made extensive use of agricultural meadows during autumn; such areas provided deer with succulent forage at a time when the nutritional quality of summer and winter range vegetation was declining because of plant senescence. Hence, Garrott et al. (1987) recommended irrigation and fertilization programs designed to retain succulent forage late into the growing season and restrictions on livestock grazing to avoid competition during periods of heavy deer use. Similar pasture management on WW herd winter ranges and holding areas might enhance deer productivity by sustaining animal condition during periods of drought.

⁹Hall, W.K. 1973. Natality and mortality of white-tailed deer in Camp Wainwright, Alberta. M.S. Thesis, University of Calgary, Alberta, Canada.

ACKNOWLEDGMENTS

I thank R. Thomas for organizing the collections and helpful comments on an early draft of the manuscript. Major assistance in deer collection was provided by R. Schaefer, J. Davis, M. Oliver, R. Mohr, R. Padgett, P. Swift, and C. Mortimore. I am grateful to V. Bleich for statistical advice. Editorial comments by V. Bleich, J. Kie, K. Whitten, and S. Holl improved the manuscript greatly. This study was funded through the California Deer Herd Management Plan Implementation Program, under Contract FG-1230.

LITERATURE CITED

- Anderson, A.E. 1981. Morphological and physiological characteristics. Pages 27-97 *in*: O.C. Wallmo, editor. Mule and black-tailed deer of North America. University of Nebraska Press, Lincoln, Nebraska, USA.
- Anderson, A.E., D.E. Medin, and D.C. Bowden. 1972. Indices of carcass fat in a Colorado mule deer population. *Journal of Wildlife Management* 36:579-594.
- Batcheler, C.L. and C.M.H. Clark. 1970. Note on kidney weights and the kidney fat index. *New Zealand Journal of Science* 13:663-668.
- Bischoff, A.I. 1958. Productivity in some California deer herds. *California Fish and Game* 44:253-259.
- Cheatum, E.L. 1949. Bone marrow as an index to malnutrition. *New York State Conservationist* 3:19-22.
- Connolly, G.E. 1981. Assessing populations. Pages 287-345 *in*: O.C. Wallmo, editor. Mule and black-tailed deer of North America. University of Nebraska Press, Lincoln, Nebraska, USA.
- Folk, M.J. and W.D. Klimstra. 1991. Reproductive performance of female key deer. *Journal of Wildlife Management* 55:386-390.
- Garrott, R.A., G.C. White, R.M. Bartmann, L.H. Carpenter, and A.W. Alldredge. 1987. Movements of female mule deer in northwest Colorado. *Journal of Wildlife Management* 51:634-643.
- Gilbert, P.F., O.C. Wallmo, and R.B. Gill. 1970. Effect of snow depth on mule deer in Middle Park, Colorado. *Journal of Wildlife Management* 33:1042-1045.
- Gill, R.B. 1972. Productivity studies of mule deer in Middle Park, Colorado. Pages 40-47 *in*: N.J. Papez, editor. *Proceedings of the Second Annual Mule Deer Workshop*, Elko, Nevada, USA.
- Hudson, P. and L.G. Browman. 1959. Embryonic and fetal development of mule deer. *Journal of Wildlife Management* 23:295-304.
- Julander, O., W.L. Robinette, and D.A. Jones. 1961. Relation of summer range condition to mule deer herd productivity. *Journal of Wildlife Management* 25:54-60.
- Kie, J.G., M. White, and D.L. Drawe. 1983. Condition parameters of white-tailed deer in Texas. *Journal of Wildlife Management* 47:583-594.
- Larson, J.S. and R.D. Taber. 1980. Criteria of sex and age. Pages 143-202 *in*: S.D. Schemnitz, editor. *Wildlife management techniques manual*. The Wildlife Society, Washington, D.C., USA.
- Leach, H.R. 1956. Food habits of Great Basin deer herds of California. *California Fish and Game* 42:243-308.
- Loft, E.R., R.C. Bertram, and D.L. Bowman. 1989. Migration patterns of mule deer in the central Sierra Nevada. *California Fish and Game* 75:11-19.

- Mayer, K.E. and W.F. Laudenslayer, Jr. 1988. A guide to wildlife habitats of California. California Department of Forestry and Fire Protection, Sacramento, California, USA.
- McCullough, D.R. 1979. The George Reserve deer herd: Population ecology of a K-selected species. University of Michigan Press, Ann Arbor, Michigan, USA.
- Morton, G.H. and E.L. Cheatum. 1946. Regional difference in breeding potential of white-tailed deer in New York. *Journal of Wildlife Management* 10:242-248.
- Ozoga, J.J. and L.J. Verme. 1982. Physical and reproductive characteristics of a supplementally-fed white-tailed deer herd. *Journal of Wildlife Management* 46:281-301.
- Pojar, T.M. and D.F. Reed. 1974. The relation of three physical condition indices of mule deer. Colorado Division of Game, Fish, and Parks, Game Information Leaflet 96:1-4.
- Ransom, A.B. 1965. Kidney and marrow fat as indicators of white-tailed deer condition. *Journal of Wildlife Management* 29:397-398.
- Ransom, A.B. 1967. Reproductive biology of white-tailed deer in Manitoba. *Journal of Wildlife Management* 31:114-123.
- Riney, T. 1955. Evaluating condition of free-ranging red deer (*Cervus elaphus*), with special reference to New Zealand. *New Zealand Journal of Science and Technology* 36:429-463.
- Robbins, C.T., A.N. Moen, and J.T. Reid. 1974. Body composition of white-tailed deer. *Journal of Animal Science* 38:871-876.
- Robinette, W.L. 1976. Mule deer mortality from various causes. Pages 125-127 in: G.W. Workman and J.B. Low, editors. Mule deer decline in the west: A symposium. Utah State University, Logan, Utah, USA.
- Robinette, W.L., J.S. Gashwiler, D.A. Jones, and H.S. Crane. 1955. Fertility of mule deer in Utah. *Journal of Wildlife Management* 19:115-136.
- Robinette, W.L., J.B. Low, and D.A. Jones. 1957. Differential mortality of sex and age among mule deer. *Journal of Wildlife Management* 21:1-16.
- Sadleir, R.M.F.S. 1987. Reproduction of female cervids. Pages 123-144 in: C.M. Wemmer, editor. Biology and management of the Cervidae. Smithsonian Institution Press, Washington, D.C., USA.
- Saltz, D., G.C. White, and R.M. Bartmann. 1992. Urinary cortisol, urea nitrogen excretion, and winter survival in mule deer fawns. *Journal of Wildlife Management* 56:640-644.
- Short, H.L. 1981. Nutrition and metabolism. Pages 99-127 in: O.C. Wallmo, editor. Mule and black-tailed deer of North America. University of Nebraska Press, Lincoln, Nebraska, USA.
- Siegel, S. 1956. Nonparametric statistics for the behavioral sciences. McGraw-Hill Book Company, New York, New York, USA.
- Swank, W.G. 1958. The mule deer in Arizona chaparral and an analysis of other important deer herds: A research and management study. Arizona Game and Fish Department Wildlife Bulletin 3:1-109.
- Taber, R.D. and R.F. Dasmann. 1958. The black-tailed deer of the chaparral: Its life history and management in the North Coast Range of California. California Department of Fish and Game, Game Bulletin 8.
- Thomas, D.C., S.J. Barry, and H.P. Kiliaan. 1989. Fetal sex ratios in caribou: Maternal age and condition effects. *Journal of Wildlife Management* 53:885-890.
- Torbit, S.C., L.H. Carpenter, D.M. Swift, and A.W. Alldredge. 1985. Differential loss of fat and protein by mule deer during winter. *Journal of Wildlife Management* 49:80-85.
- Van Vuren, D. and B.E. Coblentz. 1985. Kidney weight variation and kidney fat index: An evaluation. *Journal of Wildlife Management* 49:177-179.
- Verme, L.J. 1965. Reproduction studies of penned white-tailed deer. *Journal of Wildlife Management* 29:74-79.

- Verme, L.J. 1967. Influence of experimental diets on white-tailed deer reproduction. Transactions of the North American Wildlife and Natural Resources Conference 32:405-420.
- Wallmo, O.C., L.H. Carpenter, W.L. Regelin, R.B. Gill, and D.L. Baker. 1977. Evaluation of deer habitat on a nutritional basis. Journal of Range Management 30:122-127.

Received: 3 May 1995

Accepted: 20 November 1995

SOUTHERN RANGE EXTENSION OF THE HARLEQUIN ROCKFISH, *SEBASTES VARIEGATUS* (SCORPAENIDAE)

JAMES WILDER ORR¹ and DAVID C. BAKER²

¹Resource Assessment and Conservation Engineering Division
and

²Resource Ecology and Fisheries Management Division
Alaska Fisheries Science Center
National Marine Fisheries Service, NOAA
7600 Sand Point Way NE, Bldg. 4, BIN C15700
Seattle, Washington 98115-0070

We provide two new southern records of the harlequin rockfish, *Sebastes variegatus* Quast, 1971, that significantly extend the known southern distribution of the species. Our southernmost specimen (Fig. 1), a 253-mm standard length (SL) female, was collected by a National Marine Fisheries Service (NMFS) Fisheries Observer during commercial benthic trawling operations 95 km southwest of Newport, Oregon (44°32'N, 124°39'W), over 60 m bottom depth, on 13 September 1994. The second specimen was taken in a benthic otter trawl during the 1995 NMFS triennial survey conducted off the Washington and Oregon coasts. This 223-mm SL female (Fig. 1) was captured on 2 August 1995 off La Push, Washington (47°55'N,



Figure 1. *Sebastes variegatus*: (top) UW 22032, female, 253 mm SL, lat. 44°32'N, long. 124°39'W; (bottom) UW 027978, female, 223 mm SL, lat. 47°55'N long. 125°15'W, off La Push, Washington.

125°15'W), over 226 m bottom depth. Both Oregon (UW 22032) and Washington (UW 027978) specimens are deposited at the University of Washington Fish Collection at the School of Fisheries.

Both specimens agree closely with Quast's (1971) original description. In life, each was reddish (light yellow in preservation) with an entirely black spinous dorsal fin membrane and with prominent dark dorsal blotches that extended across the lateral line. The posterior two-thirds of the lateral line was highlighted in a broad pink-gray zone (light yellow in preservation) that interrupted the posterior dorsal blotches. In life and preservation, the anterior dorsal blotches are uninterrupted by the lateral line, which blends in with the background coloration. The specimens also possess the following counts and characters (abbreviations and methods follow Hubbs and Lagler, 1958): D XIII, 14; A III, 7; P₁, 18; pored lateral line scales, 48 (Oregon specimen), 44 (Washington); gill rakers on first arch, 14+27=41 (Oregon) and 13+27=40 (Washington); second anal spine robust and longer than third anal spine; nasal, preocular, postocular, tympanic, and parietal head spines strong; interorbital space flat; symphyseal knob moderate.

Sebastes variegatus may be confused with *S. zacentrus* (Quast 1971, Hart 1973, Eschmeyer et al. 1983, Kramer and O'Connell³ 1986, Gillespie and Saunders 1994), which is commonly collected along the Pacific coast from the Aleutian Islands to central California and ranges to San Diego (Allen and Smith 1988). Both species are reddish rockfishes with large dorsal blotches that extend ventrally from the dorsal fin to below the lateral line, and both possess the same complement of strong head spines and a prominent second anal spine that is longer and more robust than the third anal spine.

Several characters listed in the diagnosis of Quast (1971) distinguish *S. variegatus* from *S. zacentrus*: lateral line coloration, spinous dorsal fin coloration, pectoral fin ray count, and body depth. In *S. zacentrus*, unlike *S. variegatus*, the lateral line typically blends into the intersecting dorsal blotches (occasionally, the lateral line is somewhat lighter along its entire length). In *S. zacentrus*, the spinous dorsal fin membrane may have dark areas throughout the fin, typically as continuations of dorsal blotches from the body, and may have dark coloration at the tips of the dorsal spines. *Sebastes variegatus* is the only reddish rockfish found in the Northeast Pacific with the membrane of the spinous dorsal fin nearly entirely black. The pectoral fin ray count is typically 18 in *S. variegatus* (as it is in the new material), while in *S. zacentrus* the count is typically 17 (Chen 1986). Consistent with this description, in 13 specimens of *S. zacentrus* collected with the Washington specimen of *S. variegatus*, the count was 17; in one, the count was 16. The body is more slender in *S. variegatus* than *S. zacentrus*, as demonstrated also in the new material (Fig. 2).

Our Oregon record extends the verified southern range by over 500 km. Previously, the southernmost distribution had been reported as Vancouver Island,

³ Kramer, D.E. and V.M. O'Connell. 1986. Guide to northeast Pacific rockfishes genera *Sebastes* and *Sebastolobus*. Marine Advisory Bulletin 25, University of Alaska, Fairbanks, Alaska, USA.




Figure 2. Body depth, measured from dorsal fin base to pectoral fin base, versus standard length in *Sebastes zacentrus* and *S. variegatus*. WA = Washington specimen of *S. variegatus*; OR = Oregon specimen of *S. variegatus*. All specimens of *S. zacentrus* (UW 028214) examined were obtained in the August 1995 haul off La Push, Washington that also contained the 223-mm *S. variegatus*.

British Columbia (Gillespie and Saunders 1994). Allen and Smith (1988) indicated that earlier records in the NMFS database (specimens captured in 1977) from Cape Flattery, Washington, and Bodega Bay, California, were questionable. The later distribution list of adult *Sebastes* of Matarese et al. (1989: Table 29) prepared with the assistance of J. Allen (A. Matarese, NMFS, Alaska Fisheries Science Center, Seattle, Washington, pers. comm.) lists *S. variegatus* ranging south only to British Columbia. Although our new records support the validity of the Cape Flattery record, no specimens were found to support the Bodega Bay record and it must remain suspect. The original field record indicates a single adult identified as "variegatus?" among a large collection of other *Sebastes*, including *S. diploproa* (62 specimens), *S. crameri* (one specimen), and *S. babcocki* (one specimen).

ACKNOWLEDGMENTS

We thank J. Hauser, the collector of the Oregon specimen of *S. variegatus*; T.W. Pietsch and B.K. Urbain, who provided loans of the material and access to the University of Washington Fish Collection; R.R. Lauth and M.E. Wilkins, who provided access to field records and the NMFS survey database; and A.W. Kendall Jr., who critically reviewed the manuscript.

LITERATURE CITED

- Allen, M.J. and G.B. Smith. 1988. Atlas and zoogeography of common fishes in the Bering Sea and northeastern Pacific. U.S. Department of Commerce, NOAA Technical Report NMFS 66.
- Chen, L. 1986. Meristic variation in *Sebastes* (Scorpaenidae), with an analysis of character association and bilateral pattern and their significance in species separation. U.S. Department of Commerce, NOAA Technical Report NMFS 45.
- Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Co., Boston, Massachusetts, USA.
- Gillespie, G.E. and M.W. Saunders. 1994. First verified record of the shortfin mako shark, *Isurus oxyrinchus*, and second records or range extensions for three additional species, from British Columbia waters. Canadian Field-Naturalist 108:347-350.
- Hart, J.L. 1973. Pacific fishes of Canada. Fisheries Research Board of Canada, Bulletin 180.
- Hubbs, C.L. and K.F. Lagler. 1958. Fishes of the Great Lakes region. University of Michigan Press, Ann Arbor, Michigan, USA.
- Matarese, A.C., A.W. Kendall, Jr., D.M. Blood, and B.M. Vinter. 1989. Laboratory guide to the early life history stages of Northeast Pacific fishes. U.S. Department of Commerce, NOAA Technical Report NMFS 80.
- Quast, J.C. 1971. *Sebastes variegatus*, sp. n. from the northeastern Pacific Ocean (Pisces, Scorpaenidae). Fishery Bulletin 69:387-398.

Received: 1 December 1995

Accepted: 16 February 1996

DISTRIBUTION OF NESTING GREATER SANDHILL CRANES IN THE SOUTH FORK PIT RIVER VALLEY, MODOC COUNTY, CALIFORNIA

THOMAS A. ROBERTS

Environmental Science Associates
301 Brannan Street, Suite 200
San Francisco, California 94107-1849

REBECCA CULL

KEA Environmental, Inc.
455 University Avenue, Suite 370
Sacramento, California 95825

ANNE FLANNERY

Ibis Environmental Services
340 Coleman Drive
San Rafael, California 94901

The greater sandhill crane, *Grus canadensis tabida*, is a threatened species in California. Its low population is the result of habitat loss and disturbance in wintering areas in the Central Valley, summer cattle grazing in nesting areas in northeastern California (Braun et al. 1978), and high predation rates on eggs and unfledged young. Populations probably reached a low in the 1940s (Walkinshaw 1949); numbers have increased since then.

We conducted extensive sandhill crane nesting surveys in Modoc and Lassen counties in 1994 pursuant to environmental impact assessment for a major natural gas pipeline. Our survey effort included many of the areas previously surveyed in 1971, 1981, and 1988 (Littlefield¹ 1982, Littlefield² 1989). As these earlier surveys have not been replicated, we report on the part of our survey that revisited the area that Littlefield covered in the South Fork Pit River Valley. The valley is a 19-km-long, 3.2-km-wide area of mostly irrigated meadows and pastures between Highway 395 and the Westside Canal, near the town of Likely in Modoc County. The area is roughly bounded by Signal Butte on the north and by the Lassen County line on the south. This was one of the more productive areas studied by Littlefield, with 12.5% of the state's total observed greater sandhill crane nesting population in 1971 (N=112), 4.7% in 1981 (N=191), and 4.3% in 1988 (N=277).

¹ Littlefield, C.D. 1982. The status and distribution of greater sandhill cranes in California, 1981. California Department of Fish and Game, Wildlife Management Branch Administrative Report 82-1, Sacramento, California, USA.

² Littlefield, C.D. 1989. Status of greater sandhill crane breeding populations in California, 1988. California Department of Fish and Game, Wildlife Management Division, Nongame Bird and Mammal Section, Sacramento, California, USA.

We conducted preliminary ground surveys during 10-25 April 1994 to map locations of solitary birds and pairs as they began to occupy territories at the beginning of the nesting season. After consulting with C.D. Littlefield, we performed two helicopter surveys on 9 May and 3 June 1994. We flew at an altitude of 45-60 m – low enough to count individuals but not, in general, low enough to cause birds to flush. Average ground speed was 56-64 km/h, in a meandering pattern approximately 0.8 km on either side of the pipeline right-of-way (ROW) centerline, which in this area was approximately 0.4 km west of, and parallel to, Highway 395. Lateral visibility from our flight altitude gave us an effective search corridor (within which nests could clearly be seen) 1.6 km on either side of the ROW. Follow-up ground surveys confirmed numbers and locations of nesting pairs and enabled us to census all the areas of the valley where Littlefield surveyed in 1981 and 1988.

The total number of nesting pairs of greater sandhill cranes in the South Fork Pit River Valley appears to have been stable over the 24 years from 1971 to 1994, varying between 9 in 1981 and 14 in 1971 (Table 1), with a mean of 11.5 (SD = 2.08). In Littlefield's and our surveys, observers attempted a complete count of all nesting pairs in a flat, open, well-defined area, so we assume the results represent complete censuses.

Nesting use in recent years may have shifted to the northern end of the South Fork Pit River Valley (Fig. 1), possibly in response to changing land use or irrigation patterns. We documented three nests in flooded wild rice fields in the northern portion of the study area and noted that previously documented nest sites in the southern portion are heavily grazed. Another cause may be drought in the early 1990s, which reduced cover and increased predation in the southern part of the area.

Annual counts of greater sandhill crane nests in this area also are available from aerial surveys conducted by the California Department of Fish and Game (CDFG) between 1972 and 1978 (Table 1). These data are less rigorous than those previously presented because crane observations were recorded incidental to waterfowl censuses, which involved other areas between Alturas and Likely. Although the CDFG-recorded

Table 1. Nesting pairs of greater sandhill cranes in southern Modoc County, California, 1971-1994.

<u>Year</u>	<u>Complete counts, South Fork Pit River Valley</u>	<u>CDFG aerial surveys, Alturas to Likely</u>
1971	14	
1972		12
1973		9
1974		3
1975		11
1976		8
1977		10
1978		7
1981	9	
1988	12	
1994	11	
Mean (SD)	11.5 (2.08)	8.5 (2.99)




Figure 1. Greater sandhill crane nesting pair locations in 1981, 1988, and 1994, South Fork Pit River Valley, Modoc County, California. Two nest sites recorded by Littlefield in 1981 are not included because they are >3.2 km to the west of the 1994 survey corridor.

nests are probably from the same valley, the survey results are not strictly comparable. However, CDFG counts support the hypothesis of a relatively stable breeding population, varying between 3 in 1974 and 12 in 1972, with a mean of 8.5 (SD = 2.99) and no trend ($F = 0.22$, $P = 0.66$).

ACKNOWLEDGMENTS

We thank B. Deuel of CDFG for his assistance observing cranes on the first helicopter flight.

LITERATURE CITED

- Braun, C.E., K.W. Harmon, J.A. Jackson, and C.D. Littlefield. 1978. Management of national wildlife refuges in the United States: Its impacts on birds. *Wilson Bulletin* 90:309-321.
- Walkinshaw, L.H. 1949. The sandhill cranes. Cranbrook Institute of Science, Bloomfield Hills, Michigan, USA.

Received: 5 May 1995

Accepted: 22 June 1996

A RED ABALONE TAG RETURN AFTER 16 YEARS AT LIBERTY

IAN K. TANIGUCHI and PETER L. HAAKER
California Department of Fish and Game
330 Golden Shore, Suite 50
Long Beach, California 90802

A tagged red abalone, *Haliotis rufescens*, shell was given to California Fish and Game marine biologists on 26 July 1995 by R. Gutierrez, an abalone processor at the San Pedro Fish-Santa Barbara Station processing plant in Santa Barbara, California. The exact date and location of collection could not be determined because the shell was recovered from a pile of shells that represented several loads of abalone. The date of collection can only be narrowed to the week of 23-29 July 1995.

This abalone was tagged originally on 10 July 1979 at Johnson's Lee on the south side of Santa Rosa Island (33°54'N, 120°06'19"W) during a multi-year tag-recapture growth study, but it was not observed during any of the following surveys (1980-84). When tagged, it measured 147 mm long and 113 mm wide. The recovered shell measures 198 mm by 165 mm, increases of 51 mm and 52 mm over 16 yr. This growth closely fits the Von Bertalanffy growth curve for 1979-82 (Haaker et. al.,¹ in prep.) (Fig. 1).

The tag used on this abalone was a 15-mm square piece of stainless steel sheet metal with alpha-numeric characters stamped on it and a hole in one end. A length of stainless steel wire (Type: 302/304, Condition: A, Diameter: 0.8128 mm) was threaded through the hole and wrapped back around itself, leaving a wire tail (Haaker et al.² 1986). The tag was attached by passing the wire through the two most anterior completely formed respiratory pores, wrapping it back around itself several times at the base of the tag, and trimming the excess wire. The recovered tag was in good shape, with little encrustation (Fig. 2). The two respiratory pores through which the tag was threaded were sealed from the inside by the shell layers, but the external pores were incompletely filled with shell material because of the wires. Six additional respiratory pores anterior to the tagged pores were formed after tagging; two of these were closed and four were open.

When this abalone was tagged, no sign of the boring sponge, *Cliona celata californiana*, was noted on the shell. Moderate to heavy infestation of *Cliona* is now apparent on older portions of the recovered shell. *Cliona* is a yellow boring sponge that is often found living commensally on mollusks and cirripeds. The sponge

¹ Haaker, P.L., D.O. Parker, and K.C. Barsky. In Preparation. Growth of red abalone, *Haliotis rufescens*, at Johnson's Lee, Santa Rosa Island, California. California Department of Fish and Game, Marine Resources Division, Long Beach, California, USA.

² Haaker, P.L., D.O. Parker, and K.C. Henderson. 1986. Red abalone size data from Johnson's Lee, Santa Rosa Island, collected from 1978 to 1984. California Department of Fish and Game, Marine Resources Technical Report No. 53.




Figure 1. The Von Bertalanffy growth curve generated from the tag-recapture growth data for 1979-1982 ($n=657$) at Johnson's Lee, Santa Rosa Island, California (Haaker et al.,¹ in prep.)



Figure 2. Tagged red abalone shell recovered after 16 yr at large. A. Ventral view of the inside of the shell. Ruler length is 152 mm. B. Dorsal view with the tag showing in the upper left portion of the shell. C. Close-up view of the tag.

anchors itself by drilling a network of holes into the shell of its host (MacGinitie and MacGinitie 1949). *Cliona* infestation can cause structural weakening of the shell (Hansen 1970) and a heavy infestation may retard normal growth by causing the abalone to secrete additional nacre on the inside portions of the shell to repair the boring sponge damage (Ault 1985, Haaker et al.,² 1986).

The recovery of this red abalone 16 yr after it was tagged represents the longest known period of tag retention for any abalone species. The condition of the stainless steel tag and its attachment wire is evidence that this tag type is durable and persistent.

More recently, abalone tagging studies have employed number-stamped stainless steel washers, similarly attached with wires. This simplifies the manufacture of tags, as standard washers may be used, thus eliminating the time-consuming process of

cutting sheet stainless steel. The use of washers also eliminates the possibility of injury to the abalone, as well as the person tagging, by the sharp-cornered square tags.

LITERATURE CITED

- Ault, J.S. 1985. Some quantitative aspects of reproduction and growth of the red abalone, *Haliotis rufescens* Swainson. *Journal of the World Mariculture Society* 16:398-425.
- Hansen, J.C. 1970. Commensal activity as a function of age in two species of California abalones. *Veliger* 13:90-94.
- MacGinitie, G.E. and N. MacGinitie. 1949. *Natural history of marine animals*. McGraw Hill Book Co., New York, New York, USA.

Received: 22 January 1996

Accepted: 8 May 1996

FIRST FRESHWATER RECORD OF PACIFIC LAMPREY, *LAMPETRA TRIDENTATA*, FROM BAJA CALIFORNIA, MEXICO

GORGONIO RUIZ-CAMPOS and SALVADOR GONZALEZ-GUZMAN

Laboratorio de Vertebrados, Facultad de Ciencias

Universidad Autonoma de Baja California

Apdo. Postal 1653

Ensenada, Baja California 22800, Mexico

U.S. mailing address: P.O. Box 189003-064

Coronado, California 92178

An ammocoete of the Pacific lamprey, *Lampetra tridentata*, was collected on 19 February 1995 in the lower Rio Santo Domingo, Baja California, Mexico, approximately 600 m above its mouth at Bahia San Ramon, 6 km west of Vicente Guerrero (30°43'N, 116°02'W; Fig. 1). This species had not been previously reported from freshwater in Baja California (Follett 1960, Castro-Aguirre 1978, Ruiz-Campos and Contreras-Balderas 1987, Page and Burr 1991). The previous southernmost known freshwater record for this species is the Santa Ana River in southern California (Jordan and Evermann 1896, Swift et al. 1993).

The Pacific lamprey is an anadromous species distributed from Hokkaido, Japan, through the Bering Sea and Aleutian Islands (Hart 1973) to Punta Canoas, Baja California, Mexico (Hubbs 1967, Miller and Lea 1972). This species commonly spawns in the rivers of western North America from Alaska to southern California (Hubbs and Potter 1971). In southern California, this species still maintains runs in several unaltered creeks from the Carmel River south to the Santa Ana River (Swift et al. 1993). Its habitat requirements and distribution are similar to those of the anadromous rainbow trout, *Oncorhynchus mykiss*, in southern California (Swift et al. 1993). The ammocoete lacks a sucker and lives buried in the mud, feeding on micro-organisms for about 5 yr. After metamorphosis, the young adult (about 135 mm total length [TL]) migrates downstream to the sea (Hardisty and Potter 1971).

Hubbs (1967) recorded the first marine occurrence of Pacific lamprey in Baja California, 55 km southwest of Punta Canoas (28°58'N, 115°25'W). The specimen, a juvenile male 170 mm TL, was captured during a pelagic research trawl along with Pacific hake, *Merluccius productus*. Hubbs (1967:304) speculated that the juvenile lamprey "hitch-hiked a ride on a hake" from "one of the streams tributary to Monterey Bay, central California, the southernmost streams in which any massive spawning of the species has been recorded." Our specimen of Pacific lamprey from the lower Rio Santo Domingo represents the southernmost freshwater record of this species in western North America.

The specimen was captured with a minnow seine at a branch in the stream, 5 m wide, 40 cm deep, bottom of sand and gravel, and salinity 0.3 ppt, flowing into a lagoon near the mouth of Rio Santo Domingo. The ammocoete (126.5 mm TL, 3.0 g) was preserved in 10% formalin and identified by the following diagnostic




Figure 1. Map showing the geographical location of the mouth of Río Santo Domingo, Baja California, Mexico.

characteristics (Hubbs 1967, Hart 1973, Page and Burr 1991): hood surrounding toothless mouth, nasal pit, confluent gill openings, and body brown on sides and back and light silver below. Measurements (proportions of TL) and counts of the specimen are as follows: tail length, 0.289; body depth, 0.058; branchial length, 0.111; height of first dorsal (with fleshy base), 0.006; height of second dorsal (with fleshy base), 0.014; length of first dorsal, 0.153; and number of trunk myomeres, 67.

The specimen (UABC-0111) is deposited in the Fish Collection, Facultad de Ciencias, Universidad Autónoma de Baja California, at Ensenada, Baja California, Mexico.

ACKNOWLEDGMENTS

We thank A. Gerardo and V. Salceda for assistance during the field work. Also, we thank M.G. Hammann, E.P. Pister, and one anonymous reviewer for their valuable comments on the manuscript. This contribution was supported by the Consejo Nacional de Ciencia y Tecnología, Mexico (grants CONACyT 0340-N9107 and 1993P-N9507).

LITERATURE CITED

- Castro-Aguirre, J.L. 1978. Catálogo sistemático de los peces marinos que penetran en las aguas continentales de México, con aspectos zoogeográficos y ecológicos. Instituto Nacional de Pesca, México. Serie Científica Número 19:1-298.
- Follett, W.I. 1960. The freshwater fishes: Their origins and affinities. Symposium on the biogeography of Baja California and adjacent seas. Systematic Zoology 9:212-232.
- Hardisty, M.W. and I.C. Potter. 1971. The general biology of adult lampreys. Pages 127-206 in: M.W. Hardisty and I.C. Potter, editors. The biology of lampreys, volume I. Academic Press, London, England.
- Hart, J.L. 1973. Pacific fishes of Canada. Bulletin of the Fisheries Research Board of Canada 180.
- Hubbs, C.L. 1967. Occurrence of the Pacific lamprey, *Entosphenus tridentatus*, off Baja California and in streams of southern California; with remarks on its nomenclature. Transactions of the San Diego Society of Natural History 14:301-312.
- Hubbs, C.L. and I.C. Potter. 1971. Distribution, phylogeny and taxonomy. Pages 1-65 in: M.W. Hardisty and I.C. Potter, editors. The biology of lampreys, volume I. Academic Press, London, England.
- Jordan, D.S. and B.W. Evermann. 1896. The fishes of North and Middle America. Bulletin of the United States National Museum 47.
- Miller, D.J. and R.N. Lea. 1972. Guide to the coastal marine fishes of California. California Department of Fish and Game, Fish Bulletin 157.
- Page, L.M. and B.M. Burr. 1991. A field guide to freshwater fishes: North America north of Mexico. Houghton Mifflin Co., Boston., Massachusetts, USA.
- Ruiz-Campos, G. and S. Contreras-Balderas. 1987. Ecological and zoogeographical checklist of the continental fishes of the Baja California peninsula. Proceedings of the Desert Fishes Council 17:105-117.
- Swift, C.C., T.R. Haglund, M. Ruiz, and R.N. Fisher. 1993. The status and distribution of the freshwater fishes of southern California. Bulletin of the Southern California Academy of Sciences 92:101-167.

Received: 26 February 1996

Accepted: 5 April 1996

PAST OCCURRENCE OF EULACHON, *THALEICHTHYS PACIFICUS*, IN STREAMS TRIBUTARY TO HUMBOLDT BAY, CALIFORNIA

MARK R. JENNINGS¹
Department of Fisheries
School of Natural Resources
Humboldt State University
Arcata, California 95521-9905

The substantial decline of eulachon, *Thaleichthys pacificus*, in northern California over the past two decades has stimulated efforts to document past occurrences of this fish at the southern edge of its range (Moyle et al.² 1995). The most southern spawning run of eulachon was in the Mad River, Humboldt County, California (Odemar 1964). South of this drainage, eulachon have occasionally been found during the winter in Humboldt Bay (Barnhart et al. 1992), but there are no reports of adults in tributary streams. I present evidence that, in the recent past, eulachon spawned in streams tributary to Humboldt Bay.

On 10 May 1977, I found a dead, spawned-out, male eulachon (211 mm total length [TL]) on the screen of a McBane downstream migrant trap located on Jolly Giant Creek, a small stream that flows into Humboldt Bay approximately 7 km south of the Mad River. Four days later, Mr. W.G. Harper collected a second male eulachon (212 mm TL, 53 g) in spawning colors from a McBane trap on Jacoby Creek, 1.5 km south of Jolly Giant Creek. Both specimens were identified by R.A. Behrstock (formerly of Humboldt State University) who reported six other adult eulachon observed by Harper in the same McBane trap on Jacoby Creek (R.A. Behrstock, pers. comm.).

Previous records of eulachon in Humboldt Bay are rare (Emmett et al. 1991). Gotshall et al. (1980:229) reported this species as an "occasional visitor" and Samuelson³ (1973) collected one adult from South Bay on 23 April 1970. Larval stages of this fish are poorly described (Young⁴ 1984) and this may contribute to the lack of historical larval collections in the bay (e.g., see Eldridge and Bryan 1972).

¹ Current address: National Biological Service, California Science Center, Piedras Blancas Field Station, P.O. Box 70, San Simeon, California 93452-0070.

² Moyle, P.B., R.M. Yoshiyama, J.E. Williams, and E.D. Wikramanayake. 1995. Fish species of special concern in California (second edition). California Department of Fish and Game, Inland Fisheries Division, Rancho Cordova, California, USA. Final Report under Contract 2128IF.

³ Samuelson, C.E. 1973. Fishes of south Humboldt Bay, Humboldt County, California. M.S. Thesis, Humboldt State University, Arcata, California, USA.

⁴ Young, J.S. 1984. Identification of larval smelt (Osteichthyes: Salmoniformes: Osmeridae) from northern California. M.S. Thesis, Humboldt State University, Arcata, California, USA.

Three other osmerids, *Hypomesus pretiosus*, *Spirinchus starksi*, and *S. thaleichthys*, are relatively common in Humboldt Bay and their larvae have been frequently noted during past surveys (DeGeorges⁵ 1972, Emmett et al. 1991, Barnhart et al. 1992).

Spawning of eulachon in streams tributary to Humboldt Bay may be natural or the result of straying from large populations in the Mad River during the early to mid-1970s (see the accounts in Moyle et al.² 1995). Until about 2,000 years ago, the Mad River flowed into Humboldt Bay (Vick⁶ 1988) and, as recently as the 1850s, a canal connected the Mad River to Arcata Bay (Coy 1929). Such activities may have allowed the straying of eulachon into Humboldt Bay streams to produce intermittent spawning runs.

ACKNOWLEDGMENTS

This work resulted from a senior thesis in fisheries at Humboldt State University. I thank G.H. Allen, R.A. Barnhart, R.A. Behrstock, R.A. Fritzsche, W.G. Harper, and D.A. LaFaunce for their contributions to the manuscript. R.N. Lea made several useful comments on an early version of the manuscript and N.J. Scott, Jr. reviewed the penultimate draft.

LITERATURE CITED

- Barnhart, R.A., M.J. Boyd, and J.E. Pequegnat. 1992. The ecology of Humboldt Bay, California: An estuarine profile. U.S. Fish and Wildlife Service, Biological Report 1.
- Coy, O.C. 1929. The Humboldt Bay region 1850-1875. The California State Historical Association, Los Angeles, California, USA.
- Eldridge, M.B. and C.F. Bryan. 1972. Larval fish survey of Humboldt Bay. National Oceanic and Atmospheric Administration Technical Report NMFS SSRF-665.
- Emmett, R.L., S.L. Stone, S.A. Hinton, and M.E. Monaco. 1991. Distribution and abundance of fishes and invertebrates in west coast estuaries. Volume II: Species life history summaries. Estuarine Living Marine Resources Report 8. National Oceanic and Atmospheric Administration/National Ocean Service, Strategic Environmental Associates Division, Rockville, Maryland, USA.
- Gotshall, D.W., G.H. Allen, and R.A. Barnhart. 1980. An annotated checklist of fishes from Humboldt Bay, California. California Fish and Game 66:220-232.
- Odemar, M.W. 1964. Southern range extension of the eulachon, *Thaleichthys pacificus*. California Fish and Game 50:305-307.

Received: 19 March 1996

Accepted: 22 June 1996

⁵ DeGeorges, A. 1972. Feasibility of artificial reefs in intertidal waters. M.S. Thesis, Humboldt State University, Arcata, California, USA.

⁶ Vick, G.S. 1988. Late holocene paleoseismicity and relative sea level changes of the Mad River Slough, northern Humboldt Bay, California. M.S. Thesis, Humboldt State University, Arcata, California, USA.

INSTRUCTIONS FOR AUTHORS

California Fish and Game is a professional, scientific journal devoted to the conservation and understanding of California's flora and fauna. Original manuscripts dealing with California species or providing information of direct interest and benefit to California researchers and managers are welcome.

MANUSCRIPTS: Refer to the CBE Style Manual (5th Edition) and a recent issue of *California Fish and Game* for general guidance in preparing manuscripts. Specific guidelines are available from the Co-Editors-in-Chief.

COPY: Use good quality 215 x 280-mm paper. Double-space throughout with 25-mm margins. Do not hyphenate at the right margin or right-justify text. Authors should submit four good copies of their manuscript, including tables and figures, to the Co-Editors-in-Chief. An electronic copy of the manuscript on diskette in word processor format will be required with the final accepted version.

CITATIONS: All citations should follow the name-and-year system. See a recent issue of *California Fish and Game* for the format of citations and Literature Cited. Completely spell out publication and periodical names in Literature Cited. Avoid references to unpublished literature.

ABSTRACTS: Every article, except notes, must be introduced by an abstract. Abstracts should be about one typed line per typed page of text. In one paragraph describe the problem studied, most important findings, and implications of the results.

TABLES: Start each table on a separate page and double-space throughout. Do not use vertical rules. Use tabs, not the spacebar, to space between columns. Footnotes in tables should be consecutive lower-case letters, with the sequence beginning again in each table.

FIGURES: Consider proportions of figures in relation to the usable page size of *California Fish and Game* (117 x 186 mm). Figures, including captions, cannot exceed this size. Figures and line-drawings should be clear, with well-defined lines and lettering. Lettering style should be the same throughout and large enough to be readable when reduced to finished size. Type figure captions on a separate page. High-quality photographs with strong contrast are acceptable and should be submitted on glossy paper. On the back and top of each figure or photograph, lightly write the figure number and senior author's last name. Be prepared to provide high-quality, scannable, original figures or graphics files on diskette with the final accepted manuscript.

PAGE CHARGES AND REPRINTS: All authors will be charged \$40 per printed page and will be billed before publication of the manuscript. Explicit acceptance of page charges should be included in the submittal letter. Authors will receive a reprint order form along with the galley proof.

CALIFORNIA
FISH & GAME



Editor, ***CALIFORNIA FISH AND GAME***
California Department of Fish and Game
1416 Ninth St., Sacramento, CA 95814

BULK RATE
U.S. POSTAGE
PAID
Sacramento, CA.
Permit No. 949